

Pest Mole Cricket Management ¹

E. A. Buss, J. L. Capinera, and N. C. Leppla²

Three species of mole crickets were detected in the southeastern U.S. about 1900, and are now serious plant pests. Those species are the tawny mole cricket, *Scapteriscus vicinus* (Figure 1), southern mole cricket, *S. borellii* (Figure 2), and shortwinged mole cricket, *S. abbreviatus* (Figure 3). Other, non-damaging mole crickets occur in North America (e.g., *Neocurtilla hexadactyla*), but these three are the most damaging.



Figure 1. Tawny mole cricket adult. Credits: L. Buss, University of Florida



Figure 2. Southern mole cricket adult. Credits: L. Buss, University of Florida



Figure 3. Shortwinged mole cricket adult. Credits: L. Buss, University of Florida

Host Range

Though usually considered turfgrass pests, *Scapteriscus* spp. mole crickets have a broad diet. The

1. This document is ENY-324 (LH039), one of a series of the Department of Entomology and Nematology, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Date first printed October 1993. Revised: June 2006. Please visit the EDIS Website at <http://edis.ifas.ufl.edu>.

2. E. A. Buss, assistant professor, J. L. Capinera, department chairman, and N. C. Leppla, professor, Department of Entomology and Nematology, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition. All chemicals should be used in accordance with directions on the manufacturer's label. Use pesticides safely. Read and follow directions on the manufacturer's label.

southern mole cricket feeds mainly on other insects, and the tawny and shortwinged mole crickets feed on plants. Commonly injured plants include tomato, strawberry, beet, cabbage, cantaloupe, carrot, cauliflower, collard, eggplant, kale, lettuce, onion, pepper, potato, spinach, sweet potato, turnip, chufa, peanut, sugar cane, tobacco, such flowers as coleus, chrysanthemum, and gypsophila, as well as weeds such as pigweed. The tawny mole cricket often injures bahiagrass and bermudagrass, and the shortwinged mole cricket often attacks St. Augustinegrass and bermudagrass.

Damage

Mole crickets can damage plants by feeding at night on aboveground foliage or stem tissue and belowground on roots and tubers. Seedlings may be girdled at the stems near the soil surface, though some plants may be completely severed and pulled into a tunnel to be eaten. Mole cricket tunneling near the soil surface (Figures 4, 5) dislodges plants or causes them to dry out. Small mounds of soil are also pushed up. Tunneling reduces the aesthetic quality of turfgrass, interferes with the roll of the ball on golf courses, and results in reduced livestock grazing on severely infested pastures.



Figure 4. Mole cricket tunneling in dirt. Credits: E. A. Buss, University of Florida

Life Cycle

The southern and tawny mole crickets are similar in appearance and biology. The shortwinged mole cricket is different because its short wings prevent flight, and males have no calling song. Typically, the




Figure 5. Large area of mole cricket tunneling. Credits: E. A. Buss, University of Florida

eggs of these three species are laid in April-May, and nymphs occur through August. In southern Florida, however, the shortwinged mole cricket can produce eggs throughout the year. Some adults occur in August or September, but both nymphs and adults overwinter. Overwintering nymphs become adults by April, and adults fly and mate. One generation per year is normal, though in southern Florida southern mole crickets have two generations and fly three times (spring, summer, and autumn). Due to temperature differences, adult southern and tawny mole crickets emerge earlier in the year in southern Florida than in northern Florida.

Description

Eggs: About 25-60 eggs per clutch are laid in a chamber in the soil. The chamber can be 5-30 cm below the soil surface, and measures 3-4 cm in length, width, and height. Egg development lasts about 3 weeks, depending on soil temperature. Females may lay up to 5 clutches of eggs before dying.

Nymphs: Nymphs hatch from eggs from April to June. Nymphs resemble the adults, but their wings (wing pads) are not completely developed. The number of instars is variable, probably 8-10.

Adults: Mole crickets have enlarged forelegs that they use to dig in soil (Figure 6). The forelegs have large blade-like projections, called dactyls, and the number and arrangement of dactyls, and the pronotum pattern are used to identify different species (Figure 7).




Figure 6. Mole cricket anatomy.




Figure 7. Key traits used to identify *Scapteriscus* spp. mole crickets.

Tawny and southern mole cricket males attract females by producing a courtship song from their burrows early in the night. Mating and dispersal flights occur in spring, from late February to April.

Tawny mole crickets look similar to southern mole crickets, with moderately long forewings and long hind wings, a yellowish brown body, and a dark pronotum with a central band. Dactyl spacing distinguishes between the two species. The dactyls nearly touch at the base, like the shape of a "V". The tawny mole cricket's calling song occurs during the first 90 minutes after sunset.

Southern mole crickets have long hind wings that extend past the abdomen and are rounded at the tips. Adults are brownish-gray with a dark pronotum. The dactyl spacing is similar for shortwinged mole crickets (looks like a "U"), so these two species can best be distinguished by wing length. The southern

mole cricket's calling song occurs during the first two hours after sunset. As a predator, the southern mole cricket is more active and tunnels more than the tawny mole cricket.

Shortwinged mole cricket adults are 22 to 29 mm long, with wings shorter than the pronotum (area behind the head). The body is mostly tan in color, but the pronotum is brown mottled with darker spots. The top of the abdomen has a central row of large spots, and smaller spots to either side. Dactyl spacing looks like a "U". Shortwinged mole cricket males have no calling song.

Management

Sampling

Several methods are used to estimate mole cricket populations and assist in timing pesticide applications. One way is to rate the amount of tunneling damage that is visible. Tunneling is most obvious in low-cut grass or areas with minimal vegetation, and thus can be detected easily in crops, bahiagrass lawns and pastures, or bermudagrass fairways. The tunnels are most visible in early morning, when the dew is on the grass and the soil may be moist.

A more consistent but labor intensive method of sampling is a "soap flush." Flushing is more effective in moist soil. Mix 1.5 oz (2 TBSP) of lemon liquid dishwashing soap in 2 gal of water in a sprinkling can, and pour the solution onto 3-4 sq. ft. of turf. If two to four mole crickets emerge within 3 minutes after applying the soap solution, insecticide use may be justified. Flushing with a synergized pyrethrin insecticide solution is equally effective.

Adult females can also be captured at night with commercially available electronic sound traps that mimic male mating songs. In addition, adult mole crickets are strongly attracted to lights during their spring dispersal flights.

Biological Control

Mole cricket natural enemies do exist naturally in North America. Among the natural enemies are amphibians (e.g., toads, *Bufo* spp.), birds (e.g., sandhill cranes, *Grus canadensis*), and mammals

(e.g., armadillos, *Dasyopus novemcinctus*). They, and the few predatory insects that attack crickets such as tiger beetles, are not effective. Thus, several natural enemies have been introduced from South America. A parasitic wasp, *Larra bicolor* (Figure 8), was imported from South America in the 1980s and released, and has since spread in Florida. Another parasitoid, the red-eyed brazilian fly, *Ormia depleta* (Figure 9), was released in 1988. It is attracted to the mating calls of male mole crickets.



Figure 8. Mole cricket adult parasitized by *Larra bicolor*. Credits: L. Buss



Figure 9. *Ormia depleta* adult. Credits: L. Buss, University of Florida

Biological control of mole crickets can be enhanced by applying the entomopathogenic nematode, *Steinernema scapterisci* (Figure 10), or possibly other entomopathogenic nematodes. *Steinernema scapterisci* was introduced from Uruguay in 1985. It infects only *Scapteriscus* spp., persists readily in Florida's climate, and is dispersed by mole crickets. Infected mole crickets die within 2 weeks. This nematode (Nematac S®) can be purchased from commercial suppliers (Lesco, Prosource One, UHS, Gardens Alive, and others), sprayed as a suspension in water to soil, and is fairly persistent in the soil. It is more effective when

applied to adults and large nymphs in the spring or fall.



Figure 10. Mole cricket infected with the nematode, *Steinernema scapterisci*. Credits: L. Buss, University of Florida

Insecticides

Liquid and granular formulations of insecticides are commonly applied to the soil to suppress newly hatched mole cricket nymphs (Table 1) from April to June. Irrigating before an insecticide application may drive mole crickets closer to the soil surface, and helps the pesticide to penetrate into the soil. Some insecticides should be watered in after application to move them into the root zone of the plants where the mole crickets are feeding. However, it is essential to read and understand the insecticide label carefully for application directions.

Bait formulations are useful against larger nymphs in late summer. Mole crickets feed at night, so baits should be applied in the early evening. Baits are incompatible with irrigation and rainfall.

Additional information is available on the Mole Crickets web site: (<http://molecrickets.ifas.ufl.edu/mcri0001.htm>) (<http://molecrickets.ifas.ufl.edu/>) and the IPM/Biological Control web site: (<http://biocontrol.ifas.ufl.edu>).

Table 1. Insecticides labeled for professional use against mole crickets in Florida.

Active Ingredient	Florida Registered Products	Chemical Class	Signal Word
Acephate	Acephate Pro 75	Organophosphate	Caution
	Orthene Turf, Tree & Ornamental Spray		Caution
Bifenthrin	Talstar GC Flowable*, Granular*, EZ	Pyrethroid	Caution
	Talstar Lawn & Tree Flowable, PL Granular		Caution
Carbaryl	Parkway's Mole Cricket Bait	Carbamate	Caution
Clothianidin	Arena	Neonicotinoid	Caution
Cyfluthrin	Tempo 2 EC	Pyrethroid	Warning
	Tempo 20 WP, SC Ultra		Caution
Deltamethrin	DeltaGard GC*, T&O	Pyrethroid	Caution
Fipronil	Chipco Choice	Phenylpyrazole	Caution
	Top Choice		Caution
Imidacloprid	Merit 75 WP, 0.5 G	Neonicotinoid	Caution
Indoxacarb	Advion Mole Cricket Bait	Indoxacarb	Caution
Lambda-cyhalothrin	Battle GC T&O*	Pyrethroid	Caution
	Scimitar CS, GC*		Caution
Permethrin	Permethrin Pro-Termite T/O	Pyrethroid	Caution
<i>Steinernema scapterisci</i>	Nematac S	Biopesticide	Caution
Trichlorfon	Dylox 6.2 G	Organophosphate	Caution
	Dylox 80 T&O		Warning
* = Restricted Use Pesticide			

Table 2. Insecticides labeled for homeowner use against mole crickets in Florida.

Active Ingredient	Florida Registered Products	Chemical Class	Signal Word
Bifenthrin	Ortho Ortho-Klor Termite and Carpenter Ant Killer	Pyrethroid	Caution
	Scotts Turf Builder with SummerGuard		Caution
Cyfluthrin	Bayer Advanced Lawn & Garden, Multi-Insect Killer	Pyrethroid	Caution
Fipronil	Over-N-Out	Phenylpyrazole	Caution
Imidacloprid	Bayer Advanced Lawn & Garden, Season-Long Grub Control	Neonicotinoid	Caution
Lambda-cyhalothrin	Triazicide	Pyrethroid	Caution
Permethrin	Spectracide Bug Stop Multi-Purpose Insect Control	Pyrethroid	Caution
<i>Steinernema scapterisci</i>	Mole Cricket Neamtodes (Gardens Alive)	Biopesticide	Caution
Trichlorfon	Bayer Advanced 24 Hour Grub Control	Organophosphate	Caution
	Hi-Yield Dylox 6.2 Granular		