TABLE 7.—INFLUENCE OF FREQUENCY OF CUTTING UPON THE ORGANIC COMPOSITION OF DALLIS, CARIB, BAHIA AND CARPET GRASSES.

Growth Stage*	Percent Oven-dry Basis					No. of Cuttings	Intervals in 1935
	Protein	Crude Fat	Crude Fiber	Ash	N.F.E.		
Dallis Grass							
Grass	18.33	2.55	27.95	7.37	43.80	12	April 26 to Dec. 4
Early hay	16.00	2.01	32.39	6.80	42.80	6	May 16 to Oct. 30
Late hay	13.64	2.17	34.74	6.50	42.95	3	May 23 to Sept. 13
Carib Grass							
Grass	15.39	1.74	25.52	7.43	49.92	11	April 26 to Dec. 4
Early hay	9.80	1.73	33.91	5.68	48.88	3	June 11 to Oct. 17
Late hay	9.54	1.30	35.03	5.52	48.61	3	June 27 to Nov. 8
Bahia Grass							
Grass	15.54	1.84	29.07	6.84	46.71	12	April 26 to Dec. 4
Early hay	12.53	1.53	33.70	6.63	45.61	6	May 29 to Oct. 30
Late hay	11.00	1.33	34.63	6.27	46.77	3	June 7 to Sept. 30
Carpet Grass							
Grass	14.42	1.94	27.41	6.41	49.82	11	April 26 to Dec. 4
Early hay	13.58	1.76	29.50	6.37	48.79	5	May 16 to Oct. 17
Late hay	12.48	1.31	30.38	6.28	49.55	4	June 29 to Nov. 8

* See page 10 for description of these.