

?*Testudo tungia* Yeh

*Testudo tungia* Yeh 1963b, p. 224, fig. 1, pl. 1.

TYPE.—Institute of Vertebrate Paleontology and Paleoanthropology (Peking); a carapace.

TYPE LOCALITY AND HORIZON.—*Gigantopithecus* Cave, Liucheng, Kwangsi, China; Lower Pleistocene.

GEOLOGIC RANGE.—Pleistocene.

GEOGRAPHIC RANGE.—Kwangsi, China.

REMARKS.—Probably not even a testudinid; perhaps referable to the genus *Cuora*.

Genera *Inquirendae*

Genus †*Cheirogaster* Bergounioux

*Cheirogaster* Bergounioux 1935, p. 78.

GENOTYPE.—*Cheirogaster maurini* Bergounioux (by monotypy).

DEFINITION.—If correctly described, this is a monotypic genus characterized by lack of entoplastron and contact between gular and pectoral scutes.

TYPE LOCALITY AND HORIZON.—Gironde, France; Eocene.

GEOLOGIC RANGE.—Eocene.

GEOGRAPHIC RANGE.—Type locality.

REMARKS.—If correctly interpreted the type shell represents a distinct genus. The presumed distinctive characters are so unusual that a re-examination of the type material is suggested. Bergounioux (1958) correctly points out that his earlier reconstruction of the type species (1935) is greatly in error when compared with the specimen available.

*Cheirogaster* †*maurini* Bergounioux

*Cheirogaster maurini* Bergounioux 1935, p. 78.

TYPE.—M. Maurin Borbereaux collection (Gironde, France); a complete shell.

TYPE LOCALITY AND HORIZON.—Gironde, France; Ludian faunal age, Late Eocene.

GEOLOGIC RANGE.—Late Eocene.

GEOGRAPHIC RANGE.—Type locality.

*Cheirogaster* †*arrahonensis* Bergounioux

*Cheirogaster arrahonensis* Bergounioux 1957, p. 1237 (Preliminary notice).

*Cheirogaster arrahonensis* Bergounioux 1958, p. 175, figs. 15, 16, pls. 32, 33.