

LIST OF FIGURES

Figure	Page
1. Location map of the 502 profiles used in the analysis (from Hayden, et al., 1975).....	7
2. Characteristics of dimensionless beach profile $\frac{h}{h_b} = \left(\frac{x}{W}\right)^m$ for various m values (from Dean, 1977).....	8
3. Equilibrium beach profiles for sand sizes of 0.2 mm and 0.6 mm $A(D = 0.2 \text{ mm}) = 0.1 \text{ m}^{1/3}$, $A(D = 0.6 \text{ mm}) = 0.20 \text{ m}^{1/3}$	9
4. Histogram of exponent m in equation $h = Ax^m$ for 502 United States East Coast and Gulf of Mexico profiles (from Dean, 1977).....	11
5. Beach profile factor, A, vs sediment diameter, D, in relationship $h = Ax^{2/3}$ (modified from Moore, 1982).....	13
6. Profile P4 from Zenkovich (1967). A boulder coast in Eastern Kamchatka. Sand diameter: 150 mm - 300 mm. Least squares value of $A = 0.82 \text{ m}^{1/3}$ (from Moore, 1982).....	14
7. Profile P10 from Zenkovich (1967). Near the end of a spit in Western Black Sea. Whole and broken shells. $A = 0.25 \text{ m}^{1/3}$ (from Moore, 1982).....	14
8. Profile from Zenkovich (1967). Eastern Kamchatka. Mean sand diameter: 0.25 mm. Least squares value of $A = 0.07 \text{ m}^{1/3}$ (from Moore, 1982).....	15
9. Model simulation of a 0.5 meter sea level rise and beach profile response with a relatively mild sloping beach (from Moore, 1982).....	17
10. Effect of varying the sediment transport rate coefficient on cumulative erosion during the simulation of Saville's (1957) laboratory investigation of beach profile evolution for a 0.2 mm sand size (from Moore, 1982).....	18
11. Model simulation of a 0.5 meter sea level rise and beach profile response with a steep 2 meter berm (from Moore, 1982).....	19
12. Model simulation of beach nourishment for a 0.15 mm sand size (from Moore, 1982).....	20
13. Comparison of the beach profile from the model and Saville's laboratory 0.2 mm sand after 25 hours (from Moore, 1982).....	22